Noname manuscript No.
(will be inserted by the editor)

A new Improvement of Ambarzumyan's Theorem

Alp Arslan Kıraç

arXiv:1612.03622v1 [math.SP] 12 Dec 2016

Received: date / Accepted: date

Abstract We extend the classical Ambarzumyan's theorem to the quasi-periodic boundary value problems by using only a part knowledge of one spectrum. We also weaken slightly the Yurko's conditions on the first eigenvalue.

Keywords Ambarzumyan theorem · Inverse spectral theory · Hill operator · eigenvalue asymptotics

Mathematics Subject Classification (2010) 34A55 · 34B30 · 34L05 · 47E05 · 34B09

1 Preliminary Improvement

In 1929, Ambarzumyan [1] proved that if $\{(n\pi)^2 : n = 0, 1, 2 \ldots\}$ is the spectrum of the boundary value problem

$$-y''(x) + q(x)y(x) = \lambda y(x), \qquad y'(0) = y'(1) = 0 \tag{1}$$

with real potential $q \in L^1[0, 1]$, then $q = 0$ a.e. Clearly, if $q = 0$ a.e., then the eigenvalues $\lambda_n = (n\pi)^2$, $n \geq 0$.

Freiling and Yurko [6] proved that it is enough to specify only the first eigenvalue rather than the whole spectrum. More precisely, the first eigenvalue denoted by λ_0 is a mean value of the potential, that is, they proved the following Ambarzumyan-type theorem:

Theorem 1 *If* $\lambda_0 = \int_0^1 q(x)\,dx$, *then* $q = \lambda_0$ *a.e.*

A. A. Kıraç
Department of Mathematics,
Faculty of Arts and Sciences, Pamukkale University,
20070, Denizli, Turkey
E-mail: aakirac@pau.edu.tr

In [13], Yurko also provided generalizations of Theorem 1 on wide classes of self-adjoint differential operators. Some of the inverse results in [13] are in the following theorem:

Theorem 2 *(a) Let*

$$\lambda_0 = \int_0^1 q(x)\,dx$$

be the first eigenvalue of the periodic boundary value problem

$$-y''(x) + q(x)y(x) = \lambda y(x), \qquad y(1) = y(0), \quad y'(1) = y'(0), \tag{2}$$

then $q = \lambda_0$ *a.e.*

(b) Let

$$\lambda_0 = \pi^2 + \frac{2}{\alpha^2+\beta^2}\int_0^1 q(x)(\alpha sin\pi x + \beta cos\pi x)^2\,dx, \tag{3}$$

for some fixed α *and* β*, be the first eigenvalue of the anti-periodic boundary value problem*

$$-y''(x) + q(x)y(x) = \lambda y(x), \qquad y(1) = -y(0), \quad y'(1) = -y'(0). \tag{4}$$

Then

$$q = \frac{2}{\alpha^2+\beta^2}\int_0^1 q(x)(\alpha sin\pi x + \beta cos\pi x)^2\,dx \quad a.e.$$

Consider the boundary value problems $L_t(q)$ generated in the space $L^2[0,1]$ by the following differential equation and quasi-periodic boundary conditions

$$-y''(x) + q(x)y(x) = \lambda y(x), \qquad y(1) = e^{it}y(0), \quad y'(1) = e^{it}y'(0), \tag{5}$$

where $q \in L^1[0,1]$ is a real-valued function and $t \in [0,\pi)\cup[\pi,2\pi)$. The operator $L_t(q)$ is self-adjoint and the cases $t = 0$ and $t = \pi$ correspond to the periodic and anti-periodic problems, respectively. Let $\{\lambda_n(t)\}_{n\in\mathcal{Z}}$ be the eigenvalues of the operator $L_t(q)$. In the case when $q = 0$, $(2\pi n + t)^2$ for $n \in \mathcal{Z}$ is the eigenvalue of the operator $L_t(0)$ for any fixed $t \in [0, 2\pi)$ corresponding to the eigenfunction $e^{i(2\pi n+t)x}$.

The result of Ambarzumyan [1] is an exception to the general rule. In general, Borg [2] proved that one spectrum does not determine the potential. Also, Borg showed that two spectra determine it uniquely. Many generalizations of the Ambarzumyan's theorem can be found in [5,4,7,8,12,3]. But as far as we know, for the first time, we extend the classical Ambarzumyan's theorem to the quasi-periodic boundary value problems $L_t(q)$ by using only a part knowledge of one spectrum (see Theorem 4). More precisely, the aim of this paper is to weaken slightly the conditions in Theorem 1 and Theorem 2 for the first eigenvalue (see Theorem 3) and to prove a new generalization of Ambarzumyan's theorem, without using any additional conditions on the potential. We also extend Theorem 1 of the previous paper [9].

Our new uniqueness-type results read as follows:

Theorem 3 *Let $\lambda_0(t)$ be the first eigenvalue of $L_t(q)$. Then:*

(a) If $\lambda_0(t) \geq t^2 + \int_0^1 q(x)\,dx$ for $t \in [0,\pi)$, then $q = \int_0^1 q(x)\,dx$ a.e.

(b) If $\lambda_0(t) \geq (2\pi - t)^2 + \int_0^1 q(x)\,dx$ for $t \in [\pi, 2\pi)$, then $q = \int_0^1 q(x)\,dx$ a.e.

Theorem 4 *Let $\lambda_0(t)$ be the first eigenvalue of $L_t(q)$, and suppose that n_0 is a sufficiently large positive integer. Then the following two assertions hold:*

(a) If $\lambda_0(t) \geq t^2$ and $\lambda_n(t) = (2n\pi + t)^2$ for $t \in [0,\pi)$ and $n > n_0$, then $q = 0$ a.e.

(b) If $\lambda_0(t) \geq (2\pi - t)^2$ and $\lambda_n(t) = (2n\pi + t)^2$ for $t \in [\pi, 2\pi)$ and $n > n_0$, then $q = 0$ a.e.

In Theorem 3 (a), the case $t = 0$ corresponds to the periodic problem (2) and the assertion of Theorem 2 (a) holds by using $\lambda_0(0) \geq \int_0^1 q(x)\,dx$ instead of $\lambda_0(0) = \int_0^1 q(x)\,dx$. In Theorem 3 (b), the case $t = \pi$ corresponds to the anti-periodic problem (4) and the assertion of Theorem 3 (b) holds by using $\lambda_0(\pi) \geq \pi^2 + \int_0^1 q(x)\,dx$ instead of (3), namely the first eigenvalue depends only on the mean value of the potential as in Theorem 1 and Theorem 2 (a).

And, for the boundary value problem (1), the form with reduced spectrum of Ambarzumyan's theorem read as follows:

Theorem 5 *(a) If $\lambda_0 \geq \int_0^1 q(x)\,dx$, then $q = \int_0^1 q(x)\,dx$ a.e.*

(b) If $\lambda_0 \geq 0$ and $\lambda_n = (n\pi)^2$ for $n > n_0$, then $q = 0$ a.e., where n_0 is a sufficiently large positive integer.

Note that, for example, in Theorem 1 if the first eigenvalue $\lambda_0 = 0$ and $\int_0^1 q(x)\,dx = 0$, then $q = 0$ a.e. The first eigenvalue-type of Ambarzumyan's theorem, such as Theorem 1-3 and 5 (a), depends on a mean value of the potential. Hence, to prove Theorem 4 and 5 (b), without imposing an additional condition on the potential such as $\int_0^1 q(x)\,dx = 0$, we have information about the first eigenvalue with a less restrictive one and a subset of the sufficiently large eigenvalues of the spectrum.

2 Proofs

Proof of Theorem 3. (a) We show that $y = e^{itx}$ is the first eigenfunction corresponding to the first eigenvalue $\lambda_0(t)$ of the operator $L_t(q)$ for $t \in [0,\pi)$. Since the test function $y = e^{itx}$ satisfies the quasi-periodic boundary conditions in (5), by the variational principle, we get

$$t^2 + \int_0^1 q(x)\,dx \leq \lambda_0(t) \leq \frac{\int_0^1 -\bar{y}y''dx + \int_0^1 q(x)|y|^2dx}{\int_0^1 |y|^2dx} = t^2 + \int_0^1 q(x)\,dx. \quad (6)$$

This implies that $\lambda_0(t) = (t^2 + \int_0^1 q(x)\,dx)$ is the first eigenvalue corresponding to the first eigenfunction $y = e^{itx}$. Substituting this into the equation

$$-y'' + q(x)y = \lambda y,$$

we get $q = \int_0^1 q(x)\,dx$ a.e.

(b) Similarly, for $t \in [\pi, 2\pi)$, the test function $y = e^{i(-2\pi+t)x}$ is the first eigenfunction corresponding to the first eigenvalue $\lambda_0(t) = (2\pi - t)^2 + \int_0^1 q(x)\,dx$. Thus, $q = \int_0^1 q(x)\,dx$ a.e. □

Proof of Theorem 4. Note that by [10] (see also [11]), without using the assumption $\int_0^1 q(x)\,dx = 0$, the eigenvalues $\lambda_n(t)$ of the operator $L_t(q)$ for $t \neq 0, \pi$ are asymptotically located such that

$$\lambda_n(t) = (2\pi n + t)^2 + \int_0^1 q(x)\,dx + O\left(n^{-1} ln|n|\right), \tag{7}$$

for $|n| \geq n_0$, where n_0 is a sufficiently large positive integer. And, by Theorem 1.1 of [3], there is a similar asymptotic formulas for the sufficiently large eigenvalues $\lambda_n(t)$ of the operator $L_t(q)$ for $t = 0, \pi$. Thus, for all $t \in [0, \pi) \cup [\pi, 2\pi)$, if $\lambda_n(t) = (2n\pi + t)^2$ for $n > n_0$, then $\int_0^1 q(x)\,dx = 0$. From Theorem 3, $q = 0$ a.e. Thus Theorem 4 (a) and (b) are proved. □

Remark. $n < 0$ implies that $-n > 0$. In Theorem 4 (a) and (b), if we use the large eigenvalues $\lambda_{-n}(t) = (2n\pi - t)^2$ for $n > n_0$ instead of $\lambda_n(t) = (2n\pi + t)^2$, then the assertions of Theorem 4 remain valid. Moreover, for all $n > n_0$, it is enough to set either of the eigenvalues $\lambda_n(t)$, $\lambda_{-n}(t)$.

Proof of Theorem 5. (a) Arguing as in the proof of Theorem 3 (a), for $\lambda_0 \geq \int_0^1 q(x)\,dx$, the test function $y = 1$ is the first eigenfunction corresponding to the first eigenvalue $\lambda_0 = \int_0^1 q(x)\,dx$. Thus, $q = \int_0^1 q(x)\,dx$ a.e.

(b) It follows from the asymptotic formula (1.6.6) in [8] that if $\lambda_n = (n\pi)^2$ for $n > n_0$, then certainly $\int_0^1 q(x)\,dx = 0$. From (a), $q = 0$ a.e. □

References

1. Ambarzumian, V.: Über eine Frage der Eigenwerttheorie. Zeitschrift für Physik **53**, 690–695 (1929)
2. Borg, G.: Eine umkehrung der Sturm-Liouvilleschen eigenwertaufgabe bestimmung der differentialgleichung durch die eigenwerte. Acta Math. **78**, 1–96 (1946)
3. Cheng, Y.H., Wang, T.E., Wu, C.J.: A note on eigenvalue asymptotics for Hill's equation. Appl. Math. Lett. **23**(9), 1013–1015 (2010)
4. Chern, H.H., Lawb, C.K., Wang, H.J.: Corrigendum to Extension of Ambarzumyan's theorem to general boundary conditions. J. Math. Anal. Appl. **309**, 764–768 (2005)
5. Chern, H.H., Shen, C.L.: On the n-dimensional Ambarzumyan's theorem. Inverse Problems **13**(1), 15–18 (1997)
6. Freiling, G., Yurko, V.A.: Inverse SturmLiouville Problems and Their Applications. NOVA Science Publishers, New York (2001)
7. Hochstadt, H., Lieberman, B.: An inverse sturm-liouville problem with mixed given data. SIAM J. Appl. Math. **34**, 676–680 (1978)
8. Levitan, B.M., Gasymov, M.G.: Determination of a differential equation by two of its spectra. Usp. Mat. Nauk **19**, 3–63 (1964)

9. Kıraç, A.A.: On the Ambarzumyan's theorem for the quasi-periodic problem. Analysis and Mathematical Physics, **http://dx.doi.org/10.1007/s13324-015-0118-0,**, 1–4 (2015)
10. Veliev, O.A., Duman, M.: The spectral expansion for a nonself-adjoint Hill operator with a locally integrable potential. J. Math. Anal. Appl. **265**, 76–90 (2002)
11. Veliev, O.A., Kıraç, A.A.: On the nonself-adjoint differential operators with the quasiperiodic boundary conditions. International Mathematical Forum **2**(35), 1703–1715 (2007)
12. Yang, C.F., Huang, Z.Y., Yang, X.P.: Ambarzumyan's theorems for vectorial sturm-liouville systems with coupled boundary conditions. Taiwanese J. Math. **14**(4), 1429–1437 (2010)
13. Yurko, V.A.: On Ambarzumyan-type theorems. Applied Mathematics Letters **26**, 506–509 (2013)